arXiv:1211.4401v1 [math.CT] 19 Nov 2012

ON ∞-CATEGORIES

DANIEL GERIGK[1]

Abstract. A new definition for the notion of a (general) ∞-category is given.

∞-categories in $\mathbf{Set}_\Delta$

The notion of an ∞-category in Set_Δ was first defined by Boardman & Vogt ([1]), and was further developed most notably by Joyal ([3],[4],[5],[6]) and by Lurie ([7],[8]).

Definition. Denote by Cat the category of (small) categories. Denote by $\Delta \subset \mathrm{Cat}$ the full subcategory generated by the categories (associated to the linearly ordered sets) $\Delta^n := \mathbb{N}_{\leqslant n}$, for $n \in \mathbb{N}$, and call it the *category of simplices*. Let Set_Δ denote the category of (small) presheaves over Δ, and call its objects *simplicial sets*; the Yoneda embedding $\Delta \hookrightarrow \mathrm{Set}_\Delta$ will be used to identify Δ with a full subcategory of Set_Δ. $\Delta \subset \mathrm{Cat}$ is *dense*, i. e. the *nerve functor* $\mathrm{N}\colon \mathrm{Cat} \hookrightarrow \mathrm{Set}_\Delta$, which sends a category to the presheaf over Δ that it represents, is faithfully full.

Theorem. $\mathcal{C} \in \mathrm{Set}_\Delta$ is isomorphic to the nerve of a category in Cat iff for every inner horn $\Lambda^n_k \subset \Delta^n$, the induced map $\mathcal{C}(\Delta^n) \to \mathcal{C}(\Lambda^n_k)$ is bijective.

Definition. (BOARDMAN & VOGT, [1]) $\mathcal{C} \in \mathrm{Set}_\Delta$ is called an ∞-*category* if for every inner horn $\Lambda^n_k \subset \Delta^n$, the induced map $\mathcal{C}(\Delta^n) \to \mathcal{C}(\Lambda^n_k)$ is surjective. Denote by $\mathrm{Cat}^\Delta_\infty \subset \mathrm{Set}_\Delta$ the full subcategory generated by the ∞-categories.

Theorem. (JOYAL, [?]) There is a Cisinski model structure on Set_Δ, called the *Joyal model structure*, which has as fibrant objects precisely the ∞-categories.

Theorem. (CISINSKI, [?]) The Joyal model structure on Set_Δ is the Cisinski model structure whose localizer is generated by the set of spine inclusions.

∞-categories in $\mathbf{Set}_{\Delta\!\backslash}$

In the same spirit, I would like to propose the following generalization . . .

---

[1] UNIVERSITÄT BONN, GERMANY

Definition. Define the *category of multi-simplices* to be the full subcategory $\mathbb{\Delta} \subset \bigtimes_{k \geqslant 1} \Delta$ generated by those ∞-tuples over Δ with all but a finite number of components equal to Δ^0. By appending Δ^0s, every finite tuple over Δ canonically defines a corresponding multi-simplex, and when speaking of a finite tuple over Δ, I shall always mean the multi-simplex it defines. There is a canonical embedding $\Delta \hookrightarrow \mathbb{\Delta}$, sending A to (A). For $a_1, a_2, \ldots \geqslant 0$, define $\mathbb{\Delta}^{(a_1, a_2, \ldots)} := (\Delta^{a_1}, \Delta^{a_2}, \ldots)$. Call $A \in \mathbb{\Delta}$ *non-degenerate* if $A = (A_1, \ldots, A_n)$ and $A_1, \ldots, A_n \neq \Delta^0$ for some (uniquely determined) $n \geqslant 0$.

Definition. Let $\mathrm{Set}_{\mathbb{\Delta}}$ be the category of presheaves over $\mathbb{\Delta}$, and call its objects *multi-simplicial sets*; the Yoneda embedding $\mathbb{\Delta} \hookrightarrow \mathrm{Set}_{\mathbb{\Delta}}$ will be used to identify $\mathbb{\Delta}$ with a full subcategory of $\mathrm{Set}_{\mathbb{\Delta}}$.

Definition. Let $A = (\Delta^{a_1}, \Delta^{a_2}, \ldots) \in \mathbb{\Delta}$, and let $k \geqslant 1$, $a_k \geqslant 1$, $0 \leqslant i \leqslant a_k$. Define the *$(k,i)$-th face inclusion* of A to be the monomorphism $\delta^A_{(k,i)} = (\delta_1, \delta_2, \ldots)\colon B \hookrightarrow A$ in $\mathbb{\Delta}$, where for $n \neq k$, $\delta_n\colon \Delta^{a_n} \to \Delta^{a_n}$ is defined to be the respective identity, and $\delta_k\colon \Delta^{a_k - 1} \to \Delta^{a_k}$ is defined to be the i-th face inclusion of Δ^{a_k}. $\delta^A_{(k,i)}$ is called an *inner face inclusion* if $0 < i < a_k$. Define the *(k,i)-th face* of A to be the image $\operatorname{im} \delta^A_{(k,i)} \subset A$ in $\mathrm{Set}_{\mathbb{\Delta}}$. Define the *boundary* of A to be the multi-simplicial subset $\partial A \subset A$ that is the union of all the faces of A. Define the *(k,i)-th horn* of A to be the multi-simplicial subset $\Lambda^A_{(k,i)} \subset A$ that is the union of all the faces of A except the (k,i)-th face; call it an *inner horn* if the missing face is inner.

Definition. $\mathcal{C} \in \mathrm{Set}_{\mathbb{\Delta}}$ is called *globular* if for every multi-simplex $A = (A_1, A_2, \ldots)$ with $\mathrm{A}_k = \Delta^0$, $k \geqslant 1$, the map $\mathcal{C}((A_1, \ldots, A_{k-1})) \to \mathcal{C}(A)$ induced by the canonical morphism $A = (A_1, A_2, \ldots) \to (A_1, \ldots, A_{k-1})$ is bijective. Denote by $\mathbb{G} \subset \mathrm{Set}_{\mathbb{\Delta}}$ the full subcategory generated by the globular multi-simplicial sets.

Proposition. The functor glob: $\mathrm{Set}_{\mathbb{\Delta}} \to \mathbb{G}$, which sends a multi-simplicial set $\mathcal{C}$ to the largest globular multi-simplicial subset $\mathrm{glob}(\mathcal{C}) \subset \mathcal{C}$, is right adjoint to the inclusion functor $\mathbb{G} \hookrightarrow \mathrm{Set}_{\mathbb{\Delta}}$, i. e. $\mathbb{G} \subset \mathrm{Set}_{\mathbb{\Delta}}$ is a coreflective subcategory.

Definition. $\mathcal{C} \in \mathbb{G}$ is called a *strict ∞-category* if for every inner horn $\Lambda^A_{(k,i)} \subset A$ of a non-degenerate multi-simplex $A \in \mathbb{\Delta}$, the induced map $\mathcal{C}(A) \to \mathcal{C}(\Lambda^A_{(k,i)})$ is bijective.

Definition. Call $\mathcal{C} \in \mathbb{G}$ an *∞-category* if for every inner horn $\Lambda^A_{(k,i)} \subset A$ of a non-degenerate multi-simplex $A \in \mathbb{\Delta}$, the induced map $\mathcal{C}(A) \to \mathcal{C}(\Lambda^A_{(k,i)})$ is surjective. Denote by $\mathrm{Cat}^{\mathbb{\Delta}}_{\infty} \subset \mathbb{G}$ the full subcategory of ∞-categories.

Definition. Call $\mathcal{C} \in \mathrm{Cat}^{\mathbb{\Delta}}_{\infty}$ an *n-category*, for $n \geqslant 0$, if for every $\mathbb{\Delta}^{(a_1, a_2, \ldots)} \in \mathbb{\Delta}$, the map $\mathcal{C}(\mathbb{\Delta}^{(a_1, \ldots, a_n)}) \to \mathcal{C}(\mathbb{\Delta}^{(a_1, a_2, \ldots)})$ induced by $\mathbb{\Delta}^{(a_1, a_2, \ldots)} \to \mathbb{\Delta}^{(a_1, \ldots, a_n)}$ is bijective. Denote by $\mathrm{Cat}^{\mathbb{\Delta}}_{n} \subset \mathrm{Cat}^{\mathbb{\Delta}}_{\infty}$ the full subcategory of n-categories.

Definition. The canonical functor $\Delta \to \mathbb{\Delta}$ induces a functor $\mathrm{Set}_{\mathbb{\Delta}} \to \mathrm{Set}_{\Delta}$, which again induces a functor $\mathrm{Cat}_{\infty}^{\mathbb{\Delta}} \to \mathrm{Cat}_{\infty}^{\Delta}$.

I would like to postpone the details of the constructions in the next definition to a later paper . . .

Definition. It should now be possible to define $\mathbf{Cat}_{\infty}^{\mathbb{\Delta}}$, the ∞-category of ∞-categories. Its objects and (simplices of) 1-morphisms correspond to those of (the nerve of) $\mathrm{Cat}_{\infty}^{\mathbb{\Delta}}$; in general, $\mathbf{Cat}_{\infty}^{\mathbb{\Delta}}(A)$, for $A \in \mathbb{\Delta}$, may be constructed by exploiting the cartesian closedness of the topos $\mathrm{Set}_{\mathbb{\Delta}}$.
Define $\mathbf{Cat}_{n}^{\mathbb{\Delta}} \subset \mathbf{Cat}_{\infty}^{\mathbb{\Delta}}$ to be the ∞-subcategory generated by the n-categories. It should be the case that $\mathbf{Cat}_{n}^{\mathbb{\Delta}}$ is an $(n+1)$-category.

Definition. For $\mathcal{C} \in \mathrm{Cat}_{\infty}^{\mathbb{\Delta}}$, define the *$\infty$-category of k-morphisms* of $\mathcal{C}$ to be $\mathrm{Mor}^{k}(\mathcal{C}) := \mathcal{C} \circ \alpha^{k}$, where $\alpha^{1}\colon \mathbb{\Delta} \to \mathbb{\Delta}$ is the functor that sends $(A_1, A_2, \ldots)$ to $(\Delta^1, A_1, A_2, \ldots)$ - i. e. prepends Δ^1 - and similarly prepends id_{Δ^1} to a morphism. We get a functor $\mathrm{Mor}^{k}\colon \mathrm{Cat}_{\infty}^{\mathbb{\Delta}} \to \mathrm{Cat}_{\infty}^{\mathbb{\Delta}}$.

The definition of (∞, k)-categories and of k-tuply monoidal ∞-categories is straightforward.

Embedding $\mathbf{Cat}_{\infty}^{\Delta}$ in $\mathbf{Cat}_{\infty}^{\mathbb{\Delta}}$

Proposition. There is an adjunction

$$L\colon \mathrm{Set}_{\mathbb{\Delta}} \rightleftarrows \mathrm{Set}_{\Delta} : R, \quad L \dashv R,$$

such that the restriction $L|_{\mathbb{\Delta}}\colon \mathbb{\Delta} \to \mathrm{Set}_{\Delta}$ sends an object $A = (A_1, A_2, \ldots) \in \mathbb{\Delta}$ to $A_1 \times A_2 \times \ldots \in \mathrm{Set}_{\Delta}$ and a morphism $f = (f_1, f_2, \ldots)$ in $\mathbb{\Delta}$ to the morphism $f_1 \times f_2 \times \ldots$ in Set_{Δ}, and furthermore R sends an object $\mathcal{C} \in \mathrm{Set}_{\Delta}$ to $\mathrm{Mor}\,(L(-), \mathcal{C}) \in \mathrm{Set}_{\mathbb{\Delta}}$ and a morphism $f\colon \mathcal{C} \to \mathcal{D}$ in Set_{Δ} to the evident morphism in $\mathrm{Set}_{\mathbb{\Delta}}$.

Lemma. (Joyal, [6]) If a monomorphism $u\colon A \hookrightarrow B$ in Set_{Δ} is inner anodyne, then so is the monomorphism $u \times' v\colon (A \times T) \cup (B \times S) \subset B \times T$, for any monomorphism $S \hookrightarrow T$ in Set_{Δ}.

This lemma is the main ingredient in the proof of the following proposition . . .

Proposition. $L\colon \mathrm{Set}_{\mathbb{\Delta}} \to \mathrm{Set}_{\Delta}$ sends inner horn inclusions $\Lambda^{A}_{(k,i)} \subset A$ in $\mathrm{Set}_{\mathbb{\Delta}}$ to inner anodyne monomorphisms in Set_{Δ}.

Proposition. Letting $L' := L|_{\mathbb{G}}$ and $R' := \mathrm{glob} \circ R$, we obtain an adjunction

$$L'\colon \mathbb{G} \rightleftarrows \mathrm{Set}_{\Delta} : R', \quad L' \dashv R',$$

and by using the previous proposition, we see that R' induces an embedding $\mathrm{Cat}_{\infty}^{\Delta} \hookrightarrow \mathrm{Cat}_{\infty}^{\Delta\!\!\backslash\!\backslash}$.

References

[1] Boardman, Michael & Vogt, Rainer: HOMOTOPY INVARIANT ALGEBRAIC STRUCTURES ON TOPOLOGICAL SPACES. 1973.

[2] Cisinki, Denis-Charles: LES PRÉFAISCEAUX COMME MODÈLES DES TYPES D'HOMOTOPIE. 2006.

[3] Joyal, André: QUASI-CATEGORIES AND KAN COMPLEXES. 2002.

[4] Joyal, André & Tierney, Myles: QUASI-CATEGORIES VS SEGAL SPACES. 2006.

[5] Joyal, André: NOTES ON QUASI-CATEGORIES. 2008.

[6] Joyal, André: THE THEORY OF QUASI-CATEGORIES AND ITS APPLICATIONS. 2008.

[7] Lurie, Jacob: HIGHER TOPOS THEORY. 2006.

[8] Lurie, Jacob: HIGHER ALGEBRA. 2011.